

PCT/US 99/26588

U9/831-2



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

RECD 21 DEC 1999

WIPO

U9/831-2

Bescheinigung

<sup>4</sup>  
Certificate

Attestation

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixes à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

98870247.8

**PRIORITY  
DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts:  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN  
THE HAGUE, 26/11/99  
LA HAYE, LE





Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.  
Application no.  
Demande n° 98870247.8

Anmeldetag  
Date of filing  
Date de dépôt 10/11/98

Anmelder  
Applicant(s)  
Demandeur(s)  
THE PROCTER & GAMBLE COMPANY  
Cincinnati, Ohio 45202  
UNITED STATES OF AMERICA

Bezeichnung der Erfindung  
Title of the invention  
Titre de l'invention  
Bleaching compositions

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat  
State  
Pays

Tag  
Date  
Date

Aktenzeichen  
File no.  
Numero de dépôt

Internationale Patentklassifikation  
International Patent classification  
Classification internationale des brevets

/

Am Anmeldetag benannte Vertragsstaaten  
Contracting states designated at date of filing AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE  
Etats contractants désignés lors du dépôt

Bemerkungen  
Remarks  
Remarques



CM1948F

**BLEACHING COMPOSITIONS**

5

Andrea (NMN) AGOSTINI

10

Field of the invention

15

The present invention relates to cleaning compositions suitable for use in various laundry applications including hand and machine laundry methods, comprising a radical scavenger.

20

Background of the invention

25

Cleaning compositions are well known in the art. Such compositions generally comprise a number of active components that provide the cleaning function of the composition. When selecting components of the composition, it is therefore preferred that the components are stable in the composition so as not to lose cleaning effectiveness. In certain instances however, the reaction or interaction of components of the composition is inevitable. The cleaning effectiveness of the composition is reduced with premature decomposition of the oxidising agent.

30

35

Furthermore, it is believed that once decomposition of an oxidising agent is initiated, radicals provided as a result of the decomposition, initiate reactions with other components of the composition. For example, where brighteners are components of the composition, the brightener is decomposed, resulting in the loss of or decreased whiteness performance benefit, normally expected when incorporating a brightener.

The Applicant has thus now surprisingly found that this problem is solved by formulating a liquid bleaching composition according to the present invention. EP-A-558345 discloses hypochlorite-containing compositions comprising a source of strong alkalinity, a pH buffering means (silicate/carbonate) and radical scavengers. The radical scavengers of the present invention are not disclosed. Furthermore, it has been found that compositions of the present invention are stable on storage and provide improved whiteness performance in any laundry application even on ageing of the composition, i.e. when used after prolonged periods of storage, as compared to the whiteness performance delivered by the same composition under the same conditions but without any radical scavenger. More particularly, it has been found that the stability of the oxidising agent and/or the brightener (where present), are improved when the radical scavenger of the present invention is present versus when it is not.

A further advantage of the compositions of the present invention is that they are suitable for the cleaning of different types of fabrics including natural fabrics (e.g., fabrics made of cotton, viscose, linen, silk and wool), synthetic fabrics such as those made of polymeric fibers of synthetic origin as well as those made of both natural and synthetic fibers.

Yet, another advantage of the liquid bleaching compositions of the present invention is that said bleaching compositions are suitable for various laundry cleaning applications both when used in diluted conditions, e.g. as a detergent additive or a fully formulated laundry detergent composition, and when used in neat condition, e.g. as a liquid pretreater (spotter).

#### Summary of the invention

30


The present invention relates to a liquid cleaning composition comprising

- an oxidising agent,
- and a radical scavenger, which is selected from the group consisting of:

35

3


(i)



wherein S is a solubilising group ( $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ), each X, Y, Z, W, are -  
 5  $\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl groups, J is  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl group and M is H or an alkali metal, or mixtures thereof.


10

(ii)



naphtalene derivatives where in the position indicated with the numbers from 1  
 to 8 substituents are present (S, A, B, C, D, E, F, G, any substituents can  
 15 occupy any position) wherein S is a solubilising group ( $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ )  
 A, B, C, D, are  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or  
 branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl  
 groups, E, F and G can be  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with  
 R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and  
 20 secondary alkyl group and M is H or an alkali metal, or mixtures thereof.

(iii)



Where wherein S is a solubilising group ( $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ), R' is C2-20 linear or branched alkyl chain; M is as above, P and N can be  $-\text{OR}'$  or  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OCH}_3$ , or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl groups.

5

(iv) and mixtures thereof.

In a preferred embodiment the compositions of the present invention further comprise a brightener.

10

#### Detailed description of the invention

##### 15 Liquid cleaning compositions:

The compositions according to the present invention are in liquid form. Preferably, the compositions of the present invention are thickened. Thickening can be achieved by the addition of thickening components for example  
20 surfactants, more particularly anionic surfactants. Preferably the compositions are in aqueous form. More preferably, they comprise water in an amount of from 60% to 98% by weight, more preferably of from 80% to 97% and most preferably of from 85% to 97% by weight of the total aqueous liquid bleaching composition.

25

##### Oxidising agent

The oxidising agent of the present invention can be any known oxidising agent, the decomposition reaction of which produces free radicals. Suitable oxidising  
30 agent include peroxygen or hypohalite bleaching agent.

Suitable peroxygen bleaches to be used herein are hydrogen peroxide, water soluble sources thereof, or mixtures thereof. As used herein a hydrogen peroxide source refers to any compound that produces perhydroxyl ions when  
35 said compound is in contact with water. Suitable water-soluble sources of hydrogen peroxide for use herein include percarbonates, persilicates, persulphates such as monopersulfate, perborates, peroxyacids such as



diperoxydodecandioic acid (DPDA), magnesium perphthalic acid, perlauric acid, perbenzoic and alkylperbenzoic acids, hydroperoxides, aliphatic and aromatic diacyl peroxides, and mixtures thereof. Preferred peroxygen bleaches herein are hydrogen peroxide, hydroperoxide and/or diacyl peroxide. Hydrogen peroxide is the most preferred peroxygen bleach herein.

Suitable hydroperoxides for use herein are tert-butyl hydroperoxide, cumyl hydroperoxide, 2,4,4-trimethylpentyl-2-hydroperoxide, di-isopropylbenzene-monohydroperoxide, tert-amyl hydroperoxide and 2,5-dimethyl-hexane-2,5-dihydroperoxide. Such hydroperoxides have the advantage to be particularly safe to fabrics and color while delivering excellent bleaching performance when used in any laundry application.

Suitable aliphatic diacyl peroxides for use herein are dilauroyl peroxide, didecanoyl peroxide, dimyristoyl peroxide, or mixtures thereof. Suitable aromatic diacyl peroxide for use herein is for example benzoyl peroxide. Such diacyl peroxides have the advantage to be particularly safe to fabrics and color while delivering excellent bleaching performance when used in any laundry application.

Preferably the oxidising agent of the invention is a hypohalite bleach. Hypohalite bleaches may be provided by a variety of sources, including bleaches that are oxidative bleaches and subsequently lead to the formation of positive halide ions as well as bleaches that are organic based sources of halides such as chloroisocyanurates.

Suitable hypohalite bleaches for use herein include the alkali metal and alkaline earth metal hypochlorites, hypobromites, hypoiodites, chlorinated trisodium phosphate dodecahydrates, potassium and sodium dichloroisocyanurates, potassium and sodium trichlorocyanurates, N-chloroimides, N-chloroamides, N-chloroamines and chlorohydantoins.


Particularly preferred oxidising agents of the present invention are the alkali metal and/or alkaline earth metal hypochlorites selected from the group consisting of sodium, potassium, magnesium, lithium and calcium hypochlorites, and mixtures thereof. Most preferably the oxidising agent is sodium hypochlorite.

Preferably, the liquid compositions according to the present invention comprise said oxidising agent such that the content of active oxidising agent in the composition is from 0.1% to 20% by weight, more preferably from 0.25% to 8%  
5 by weight, most preferably from 0.5% to 6% by weight of the composition.

### The radical scavenger


The compositions of the present invention comprise a radical scavenger. The  
10 radical scavengers described herein have not previously been described in the art. The radical scavenger may be selected from one of the three classes of radical scavenger described herein or may be a combination of more than one of the radical scavengers.

15 The first class of radical scavengers has the general formula:



20 wherein S is a solubilising group ( for example  $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ), each of X, Y, Z, W may be selected from  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or branched alkyl chain C1-C20), or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl groups; J may be selected from  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-$   
25  $\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or branched alkyl chain C1-C20), or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl group; and M is H or an alkali metal, or mixtures thereof.

The second class of radical scavengers has general formula:



naphtalene derivatives where in the position of the naphthlene ring structure are given the numbers from 1 to 8 (as seen above) and each position may be substituted with a substituent groups selected from S, A, B, C, D, E, F, G wherein S is a solubilising group (for example  $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ); A, B, C, and D are selected from  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with  $\text{R}=\text{linear or branched alkyl chain C1-C20}$ ), or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl groups; E, F and G are selected from  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with  $\text{R}=\text{linear or branched alkyl chain C1-C20}$ ), or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl group; and M is selected from H or an alkali metal.

Other suitable radical scavengers are polymers where structures of Class 1 are present in a polymer, i.e. wherein S is a solubilising group ( $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ), each Y, Z, J are  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with  $\text{R}=\text{linear or branched alkyl chain C1-C20}$ ), or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl groups, X, W are two polymer chains of an homo or a co-polymer, where the unit described is repeated. is  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with  $\text{R}=\text{linear or branched alkyl chain C1-C20}$ ), or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl group and M is H or an alkali metal, or mixtures thereof.

Other suitable radical scavengers are polymers where structures 2 is present in a polymer, i.e. wherein wherein S is a solubilising group ( $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ), each C, D, E, F, G are  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with  $\text{R}=\text{linear or branched alkyl chain C1-C20}$ ), or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl groups, A, B are two polymer chains of an homo or a co-polymer, where the unit described is repeated. is  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with  $\text{R}=\text{linear or branched alkyl chain C1-C20}$ ), or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl group and M is H or an alkali metal, or mixtures thereof.

Naturally, for the purpose of the present invention, the radical scavengers have to be stable to the hypohalite bleach.

5 The compositions of the present invention comprise from 0.001% to 10% by weight of the total composition of a radical scavenger, or a mixture thereof, preferably from 0.01% to 8%, more preferably from 0.1% to 6% and most preferably from 0.2% to 4%.

10 Alternatively, the compositions of the present invention may comprise a mixture of the radical scavengers or a combination of at least one radical scavenger of the present invention and another radical scavenger known in the art or those described in the Applicants co-pending application number 97870139.9-2108 (attorney docket number CM1603).

15 Advantageously, the compositions of the invention are stable on storage. By "stable on storage", it is to be understood that the compositions of the present invention typically do not undergo more than 25% loss of available oxidising agent after 5 days of storage at  $50^{\circ}\text{C} \pm 0.5^{\circ}\text{C}$  and more preferably not more  
20 than 20%. The % loss of available chlorine may be measured using the method described, for instance, in "Analyses des Eaux et Extraits de Javel" by "La chambre syndicale nationale de L'eau de Javel et des produits connexes", pages 9-10 (1984). Said method consists in measuring the available oxidising agent in the fresh compositions, i.e. just after they are made, and in the same  
25 compositions after 5 days at  $50^{\circ}\text{C}$ .

#### The brightener

An preferred optimal component of the invention is a brightener. Any brightener  
30 known in the art may be used herein including both hydrophobic and hydrophilic brighteners and mixtures thereof.

Brighteners are compounds which have the ability to fluoresce by absorbing ultraviolet wave-lengths of light and re-emitting visible light. Brighteners, also  
35 referred to as fluorescent whitening agents (FWA), have been extensively described in the art, see for instance EP-A-0 265 041, EP-A-0 322 564, EP-A-0

317 979 or "Fluorescent whitening agents" by A.K. Sarkar, published by MERROW, especially page 71-72.

Commercial optical brighteners which may be useful in the present invention can be classified into subgroups, which include, but are not necessarily limited to, derivatives of stilbene, pyrazoline, coumarin, carboxylic acid, methinecyanine, dibenzothiophene-5,5-dioxide, azole, 5- and 6-membered-ring heterocycle, and other miscellaneous agents. Examples of such brighteners are disclosed in "The Production and Application of Fluorescent Brightening Agents", M. Zahradnik, Published by John Wiley & Sons, New York (1982). Further optical brighteners which may also be used in the present invention include naphthlimide, benzoxazole, benzofuran, benzimidazole and any mixtures thereof. Particularly preferred brighteners for use herein are the derivatives of stilbene and mixtures thereof.


Examples of optical brighteners which are useful in the present compositions are those identified in U.S. Patent 4,790,856. These brighteners include the PHORWHITE® series of brighteners from Verona. Other brighteners disclosed in this reference include: Tinopal-UNPA®, Tinopal CBS® and Tinopal 5BM® available from Ciba-Geigy; Artic White CC® and Artic White CWD®; the 2-(4-styryl-phenyl)-2H-naphtho[1,2-d]triazoles; 4,4'-bis(1,2,3-triazol-2-yl)-stilbenes; 4,4'-bis(styryl)bisphenyls; and the aminocoumarins.

Specific examples of brighteners useful herein include 4-methyl-7-diethyl-amino coumarin; 1,2-bis(-benzimidazol-2-yl)ethylene; 1,3-diphenyl-pyrazolines; 2,5-bis(benzoxazol-2-yl)thiophene; 2-styryl-naphtho-[1,2-d]oxazole; 2-(stilbene-4-yl)-2H-naphtho[1,2-d]triazole, 3-phenyl-7-(isoindolinyl) coumarin; 3-methyl-7-(isoindolinyl) coumarin; 3-chloro-7-(isoindolinyl) coumarin; 4-(isoindolinyl)-4'-methylstilbene; 4-(isoindolinyl)-4'-methoxystilbene; sodium 4-(isoindolinyl)-4'-stilbenesulfonate; 4-(isoindolinyl)-4'-phenylstilbene; 4-(isoindolinyl)-3-methoxy-4'-methylstilbene; 4-(2-chloroisoindolinyl)-4'-(2-methylisoindolinyl)-2,2'-stilbenedisulfonic acid; disodium 4,4'-diisoindolinyl-2,2'-stilbene disulfonate; 4,4'-diisoindolinyl-2,2'-stilbenedisulfonamide; disodium 4,4'-(7,8-dichloro-1-isoindolinyl)2,2-stilbenedisulfonate; disodium 4,4'-(7-chloro-1-isoindolinyl)2,2-stilbenedisulfonate; disodium 4,4'-(6-isopropoxy-1-isoindolinyl)2,2-stilbenedisulfonate; disodium 4,4'-(7,8-diisopropyl-1-isoindolinyl)2,2-stilbenedisulfonate; disodium 4,4'-(7-butoxy-1-isoindolinyl)2,2-

stilbenedisulfonate; disodium 4,4'-(6-trifluoromethyl-1-isoindolinyl)2,2-  
 stilbenedisulfonate; disodium 4,4'-[6-(1,4,7-trioxanonyl)-1-isoindolinyl)]2,2-  
 stilbenedisulfonate; disodium 4,4'-(7-methoxymethyl-1-isoindolinyl)2,2-  
 stilbenedisulfonate; disodium 4,4'-(6-phenyl-1-isoindolinyl)2,2-  
 5 stilbenedisulfonate; disodium 4,4'-(6-naphthyl-1-isoindolinyl)2,2-  
 stilbenedisulfonate; disodium 4,4'-(6-methylsulfonyl-1-isoindolinyl)2,2-  
 stilbenedisulfonate; disodium 4,4'-(7-cyano-1-isoindolinyl)2,2-  
 stilbenedisulfonate; and disodium 4,4'-[7-(1,2,3-trihydroxypropyl)-1-  
 isoindolinyl)]2,2-stilbenedisulfonate; disodium 4-isoindolinyl-4'-ethoxy-2,2'-  
 10 stilbenedisulfonate; disodium 4-isoindolinyl-4'-methoxy-2,2'-stilbenedisulfonate;  
 disodium 4-isoindolinyl-4'-ethoxy-2,2'-stilbenedisulfonamide; disodium 4-  
 isoindolinyl-4'-methyl-2,2'-stilbenedisulfonamide; disodium 4,4'-bis-(2-  
 diethanolamino-4-anilino-s-triazin-6-ylamino)stilbene-2:2 disulphonate,  
 disodium 4,4'-bis-(2-morpholino-4-anilino-s-triazin-6-ylamino)-stilbene-2:2'-  
 15 disulphonate, disodium 4,4'-bis-(2,4-dianilino -s-triazin-6-ylamino)-stilbene-2-  
 sulphonate, monosodium 4,4''-bis-(2,4-dianilino -s-triazin-6-ylamino)-stilbene-  
 2:2'-disulphonate, disodium 4,4'-bis-(2-anilino-4-(N-methyl-N-2-  
 hydroxyethylamino)-s-triazin-6-ylamino)stilbene-2,2'-disulphonate, disodium  
 4,4'-bis-(4-phenyl-2,1,3-triazol-2-yl)-stilbene-2,2' disulphonate, disodium 4,4'-  
 20 bis-(2-anilino-4-(1-methyl-2-hydroxyethylamino)-s-triazin-6-ylamino)-stilbene-  
 2,2'-disulphonate, sodium 2-(stilbyl-4''-(naphtho-1',2':4,5)-1,2,3-triazole-2''-  
 sulphonate, 4,4'-bis-(2-sulphostyryl)-biphenyl, 4,4'-bis(4-phenyl-2H-1,2,3-triazol-  
 2-yl)-2,2'-stilbenedisulfonic acid and mixture thereof. See also U.S. Patent  
 3,646,015, U.S. Patent 3,346,502 and U.S. Patent 3,393,153 for further  
 25 examples of brighteners useful herein.

Indeed one of the functionally equivalent derivative salts of 4,4'-bis(4-phenyl-  
 2H-1,2,3-triazol-2-yl)-2,2'-stilbenedisulfonic acid, namely its sodium salt is  
 available from Mobay Chemical Corporation, a subsidiary of Bayer AG under  
 30 the name Phorwite® CAN. The amine salt is available from Molay under the  
 name Phorwite® CL solution. The potassium salt is available under the name  
 Phorwite® BHC 766.

Specific examples of hydrophilic optical brighteners useful in the present  
 35 invention are those having the structural formula:



wherein  $R_1$  is selected from anilino, N-2-bis-hydroxyethyl and NH-2-hydroxyethyl;  $R_2$  is selected from N-2-bis-hydroxyethyl, N-2-hydroxyethyl-N-methylamino, morphilino, chloro and amino; and M is a salt-forming cation such as sodium or potassium.

When in the above formula,  $R_1$  is anilino,  $R_2$  is N-2-bis-hydroxyethyl and M is a cation such as sodium, the brightener is 4,4',-bis[(4-anilino-6-(N-2-bis-hydroxyethyl)-s-triazine-2-yl)amino]-2,2'-stilbenedisulfonic acid and disodium salt. This particular brightener species is commercially marketed under the tradename Tinopal-UNPA-GX by Ciba-Geigy Corporation. Tinopal-UNPA-GX is the preferred hydrophilic optical brightener useful in the detergent compositions herein.

When in the above formula,  $R_1$  is anilino,  $R_2$  is N-2-hydroxyethyl-N-2-methylamino and M is a cation such as sodium, the brightener is 4,4'-bis[(4-anilino-6-(N-2-hydroxyethyl-N-methylamino)-s-triazine-2-yl)amino]-2,2'-stilbenedisulfonic acid disodium salt. This particular brightener species is commercially marketed under the tradename Tinopal 5BM-GX by Ciba-Geigy Corporation.

When in the above formula,  $R_1$  is anilino,  $R_2$  is morphilino and M is a cation such as sodium, the brightener is 4,4'-bis[(4-anilino-6-morphilino-s-triazine-2-yl)amino]-2,2'-stilbenedisulfonic acid, sodium salt. This particular brightener species is commercially marketed under the tradename Tinopal AMS-GX by Ciba Geigy Corporation.

Other substituted stilbene 2,2'-disulfonic acid derivatives also include 4-4'-bis(2-2' styryl sulfonate) biphenyl, commercially available from Ciba-Geigy under the trade name Brightener 49® or other hydrophilic brighteners like for example Brightener 3® or Brightener 47®, also commercially available from Ciba-Geigy.

Specific examples of hydrophobic brighteners useful in the present invention include the polycyclic oxazole derivatives such as benzo-oxazole derivatives, or mixtures thereof and particularly preferred herein the benzo-oxazole derivatives. An example of such a brightener is benzoxazole,2,2'-(thiophenaldyl)bis having the following formula  $C_{18}H_{10}N_2O_2S$ , commercially available from Ciba-Geigy under the trade name Tinopal SOP®. This brightener is almost insoluble in water, i.e. it has a solubility being lower than 1 gram per liter. Another example of such a brightener is bis(sulfobenzofuranyl)biphenyl, commercially available from Ciba-Geigy under the trade name Tinopal PLC®.

10

By "hydrophobic brighteners", it is to be understood herein any brightener whose solubility in water is lower than 10 grams per liter at 25°C. By "solubility" of a given compound, it is to be understood herein the amount of said compound solubilized in deionized water at 25°C. Thus, a compound having a solubility being lower than 10 grams per liter means that when less than 10 grams of said given compound is incorporated in deionized water at 25°C said compound is entirely dissolved in said water, i.e. a clear and stable solution is obtained. In other words, incorporating 10 grams per liter or more of said given compound in water will result in a precipitation of said compound in said medium. Accordingly, by "hydrophilic brighteners", it is to be understood herein any brightener whose solubility in water is higher or equal to 10 grams per liter at 25°C.

25

Where present, brightener is incorporated at a level of from 0.001% to 1.0%, preferably from 0.005% to 0.5%, more preferably from 0.005% to 0.3% and most preferably from 0.008% to 0.1%, by weight of the composition.

30

Where hydrophobic brighteners are present in the compositions herein they may both be solubilized or suspended in the compositions of the present invention. Such brighteners solubilisation can be for example achieved by means of a surfactant or a mixture thereof as described herein after. Various surfactants may be used for this purpose like C8-C20 alkyl aryl sulphonates as described for example in U.S. Patent 4, 623,476 or amine oxides as described for example in EP-A-186386. Preferred surfactants also called "co-surfactants" to solubilise and/or suspend such a hydrophobic brightener are anionic surfactants including alkyl sulphates or alkylalkoxy sulphates having from 4 to 30 carbon atoms in the alkyl chain, or alkylethoxycarboxylates having from 6 to

35



30 carbon atoms in the alkyl chain such as AkypoSoft® 100 NV from Chemy or Sandosan LNCS from Sandoz. Preferred are C12-C14 alkyethoxysulphates. Such co-surfactants herein should be used in amounts required to solubilize the hydrophobic brightener in need thereof.

5

Generally, when a co-surfactant is used, the liquid compositions of the present inventions are prepared in a process wherein the hydrophobic brightener and the co-surfactant are first mixed to form a premix, before the premix is then mixed with the remainder of the composition which has been separately prepared.

10

Alternatively, the hydrophobic brightener may be suspended by means of a specific suspending agent, like polymers and/or colloidal particulate silicate. Any polymers known to those skilled in the art as having suspending properties are suitable for use herein including those described for example in EP-A-206718.

15

#### Optional chelating agents

20 In the compositions of the present invention, the presence of a chelating agent in addition to radical scavenger is not compulsory, but is highly preferred.

It is believed that the presence of a chelating agent in the compositions of the present invention further contributes to the brightener stability upon prolonged periods of storage, and thus to the effective whiteness performance of these compositions upon ageing.

25

Suitable chelating agents may be any of those known to those skilled in the art such as the ones selected from the group comprising phosphonate chelating agents, phosphate chelating agents, polyfunctionally-substituted aromatic chelating agents, ethylenediamine N,N'-disuccinic acids, or mixtures thereof.

30


Chelating agents may be desired in the compositions of the present invention, preferably phosphate chelating agents like phytic acid, as they further contribute to the benefit delivered by the radical scavengers herein by further improving the stability of the brighteners, thus delivering effective whiteness

35


performance in any laundry application upon ageing of the compositions, i.e. after prolonged periods of storage.

Suitable phosphonate chelating agents for use herein may include alkali metal ethane 1-hydroxy diphosphonates (HEDP), alkylene poly (alkylene phosphonate), as well as amino phosphonate compounds, including amino aminotri(methylene phosphonic acid) (ATMP), nitrilo trimethylene phosphonates (NTP), ethylene diamine tetra methylene phosphonates, and diethylene triamine penta methylene phosphonates (DTPMP). The phosphonate compounds may be present either in their acid form or as salts of different cations on some or all of their acid functionalities. Preferred phosphonate chelating agents to be used herein are diethylene triamine penta methylene phosphonate (DTPMP) and ethane 1-hydroxy diphosphonate (HEDP). Such phosphonate chelating agents are commercially available from Monsanto under the trade name DEQUEST®.

Suitable phosphate chelating agents are as following: Phosphonic acid can be condensed in the reaction :




The reaction can be repeated with any of the reactive OH groups, and phosphate ingredients are obtained which can be - linear or branched polyphosphates of structure



15


-when R is M or



5


- when M is a counterion, preferably alkali metal;
- when  $0 \leq n+m < 500$  (if  $n+m = 0$  then the compound is phosphonic acid)

10 - cyclic polyphosphates (also referred to as metaphosphates), of structure




15

-when R is M or



-if R is



- 5 the phosphate compound contains both cycles and branched chain, and can be referred to as an ultraphosphate.)  
 - where M is a counterion, preferably an alkali metal  
 - where  $0 \leq n+m < 500$
- 10 All such phosphate ingredients are suitable for use herein, and preferred are linear phosphate ingredients (i.e. R is M) where n is 1 (pyrophosphate) and n is 2 (tripolyphosphate (STPP)), most preferably where n is 2. The most commonly available form of these phosphate is where M is Sodium.
- 15 Phytic acid, which is particularly suitable for use herein, is a hexa-phosphoric acid that occurs naturally in the seeds of many cereal grains, generally in the form of the insoluble calcium-magnesium salt. It may also be derived from corn steep liquor. Commercial grade phytic acid is commercially available from J.T.Baker Co., e.g. as a 40% aqueous solution. It is intended that the present
- 20 invention covers the acidic form of phytic acid as well as alkali metal salt derivatives thereof, particularly sodium or potassium salts thereof. Sodium phytate is available from Jonas Chemical Co (Brooklyn, N.Y.). In fact since the typical pH of the compositions of the present invention are in the alkaline pH range, the phytic acid component exists primarily as the ionized salt in the liquid
- 25 compositions herein even if it is added in its acidic form. Mixtures of such salts of phytic acid are also covered.

Polyfunctionally-substituted aromatic chelating agents may also be useful in the compositions herein. See U.S. patent 3,812,044, issued May 21, 1974, to Connor et al. Preferred compounds of this type in acid form are

30 dihydroxydisulfobenzenes such as 1,2-dihydroxy -3,5-disulfobenzene.

A preferred biodegradable chelating agent for use herein is ethylene diamine N,N'- disuccinic acid, or alkali metal, or alkaline earth, ammonium or substitutes ammonium salts thereof or mixtures thereof. Ethylenediamine N,N'- disuccinic acids, especially the (S,S) isomer have been extensively described in US  
5 patent 4, 704, 233, November 3, 1987, to Hartman and Perkins. Ethylenediamine N,N'- disuccinic acids is, for instance, commercially available under the tradename ssEDDS® from Palmer Research Laboratories.

10 Particularly preferred chelating agents for use herein are phosphate chelating agents like sodium pyrophosphate, sodium tripolyphosphate, phytic acid, and mixtures thereof.

Typically, the compositions according to the present invention comprise up to  
15 10%, preferably from 0.01% to 5% by weight, more preferably from 0.05% to 2% and most preferably from 0.1% to 1.5% by weight of the composition.

#### pH

20 The pH of the liquid compositions according to the present invention, as is, is typically from 12 to 14 measured at 25°C. The liquid compositions of the invention have a pH of from 7.5 to 13, preferably from 8 to 12, more preferably from 8.5 to 11.5, when diluted into 1 to 500 times its weight of water. It is in this alkaline range that the optimum stability and performance of the hypohalite as  
25 well as fabric whiteness and/or safety are obtained. The pH range is suitably provided by a buffering component.

Suitable sources of alkalinity are the caustic alkalis such as sodium hydroxide, potassium hydroxide and/or lithium hydroxide, and/or the alkali metal oxides  
30 such as sodium and/or potassium oxide. A preferred strong source of alkalinity is a caustic alkali, more preferably sodium hydroxide and/or potassium hydroxide. Typical levels of such caustic alkalis, when present, are of from 0.1% to 1.5% by weight, preferably from 0.5% to 1.5% by weight of the composition.

35

Optional pH buffering components

In the compositions of the present invention, the presence of a pH buffering component is not compulsory, but is preferred.

5

It is believed that the presence of a pH buffering component in the compositions of the present invention further contributes to the effective whiteness performance of these compositions as well as to the fabric safety. Indeed, the pH buffering component allows to control the alkalinity in the bleaching solution, i.e. maintain the pH of the bleaching solution at a pH of at least 7.5, preferably at least 8.5, and more preferably at least 9.5 for a longer period of time starting from the moment at which the dilution is completed (e.g. when the bleaching composition of the present invention is diluted in the bleaching solution at a dilution level of 200:1 (water:composition)). It is believed that the pH buffering component may also contribute to brightener stability, where present, and thus to the effective whiteness performance of the compositions herein.

10

15

20

The pH buffering component ensures that the pH of the composition is buffered to a pH value ranging from 7.5 to 13, preferably from 8 to 12, more preferably from 8.5 to 11.5 after the composition has been diluted into 1 to 500 times its weight of water.

25

Suitable pH buffering components for use herein are selected from the group consisting of alkali metal salts of carbonates, polycarbonates, sesquicarbonates, silicates, polysilicates, boron salts, phosphates, stannates, alluminates and mixtures thereof. The preferred alkali metal salts for use herein are sodium and potassium.

30

35

Suitable boron salts or mixtures thereof for use herein include alkali metal salts of borates and alkyl borates and mixtures thereof. Examples of boron salts include boric acid, alkali metal salts of metaborate, tetraborate, octaborate, pentaborate, dodecaboron, borontrifluoride and alkyl borate containing from 1 to 12 carbon atoms, preferably from 1 to 4. Suitable alkyl borate includes methyl borate, ethyl borate and propyl borate. Particularly preferred boron salts herein are the alkali metal salts of metaborate, such as sodium metaborate, potassium metaborate, and the alkali metal salts of borate, such as sodium

borate, or mixtures thereof. Boron salts like sodium metaborate and sodium tetraborate are commercially available from Borax and Societa Chimica Larderello under the name sodium metaborate and Borax®.

- 5 Particularly preferred pH buffering components are selected from the group consisting of sodium carbonate, sodium silicate, sodium borate, sodium metaborate and mixtures thereof.

10 The raw materials involved in the preparation of hypohalite bleaches usually contain by-products, e.g calcium carbonate resulting in an amount of up to 0.4% by weight of by-product within the hypohalite composition. However, at such amount, the by-product will not have the buffering action defined above.

15 Liquid bleaching compositions herein will contain an amount of pH buffering component of from 0.5% to 9% by weight, preferably from 0.5% to 5% by weight, and more preferably in an amount of from 0.6% to 3% by weight of the composition.

20 The composition according to the invention may comprise other optional components such as bleach-stable surfactants, organic or inorganic alkalis, builders, thickening agents, polymers, pigments, dyes, solvents, perfumes, and mixtures thereof.

#### Optional surfactants

25 The liquid compositions of the present invention may further comprise surfactant as a desirable optional ingredient. Said surfactants may be present in the compositions according to the present invention in amounts of from 0.1% to 50%, preferably from 0.1% to 40% and more preferably from 1% to 30% by weight of the composition.

30 Surfactants suitable for use herein include anionic surfactants, nonionic surfactants, cationic surfactants, amphoteric surfactants, zwitterionic surfactants, and mixtures thereof.

35 Suitable anionic surfactants for use in the compositions herein include water-soluble salts or acids of the formula  $\text{ROSO}_3\text{M}$  wherein R preferably is a  $\text{C}_{10}$ -

- C<sub>24</sub> hydrocarbyl, preferably linear or branched alkyl having a C<sub>10</sub>-C<sub>20</sub> alkyl component, more preferably a C<sub>12</sub>-C<sub>18</sub> linear or branched alkyl, and M is H or a cation, e.g., an alkali metal cation (e.g., sodium, potassium, lithium), or ammonium or substituted ammonium (e.g., methyl-, dimethyl-, and trimethyl-  
5 ammonium cations and quaternary ammonium cations, such as tetramethyl-ammonium and dimethyl piperdinium cations and quaternary ammonium cations derived from alkylamines such as ethylamine, diethylamine, triethylamine, and mixtures thereof, and the like).
- 10 Other suitable anionic surfactants for use herein are water-soluble salts or acids of the formula RO(A)<sub>m</sub>SO<sub>3</sub>M wherein R is an unsubstituted C<sub>6</sub>-C<sub>24</sub> linear or branched alkyl group having a C<sub>6</sub>-C<sub>24</sub> alkyl component, preferably a C<sub>10</sub>-C<sub>20</sub> linear or branched alkyl, more preferably C<sub>12</sub>-C<sub>18</sub> linear or branched alkyl, A is  
15 an ethoxy or propoxy unit, m is greater than zero, typically between about 0.5 and about 30, more preferably between about 0.5 and about 5, and M is H or a cation which can be, for example, a metal cation (e.g., sodium, potassium, lithium, calcium, magnesium, etc.), ammonium or substituted-ammonium cation. Alkyl ethoxylated sulfates as well as alkyl propoxylated sulfates are  
20 contemplated herein. Specific examples of substituted ammonium cations include methyl-, dimethyl-, trimethyl-ammonium and quaternary ammonium cations, such as tetramethyl-ammonium, dimethyl piperdinium and cations derived from alkanolamines such as ethylamine, diethylamine, triethylamine, mixtures thereof, and the like. Exemplary surfactants are C<sub>12</sub>-C<sub>18</sub> alkyl  
25 polyethoxylate (1.0) sulfate, C<sub>12</sub>-C<sub>18</sub>E(1.0) sulfate, C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (2.25) sulfate, C<sub>12</sub>-C<sub>18</sub>E(2.25) sulfate, C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (3.0) sulfate, C<sub>12</sub>-C<sub>18</sub>E(3.0) sulfate, and C<sub>12</sub>-C<sub>18</sub> alkyl polyethoxylate (4.0) sulfate, C<sub>12</sub>-C<sub>18</sub>E(4.0) sulfate, wherein the counterion is conveniently selected from sodium and potassium.
- 30 In a particularly preferred embodiment of the present invention the surfactant system used is an anionic surfactant system, namely alkyl alkoxy (preferably ethoxy) sulphate alone for example alkyl (ethoxy) 3 sulphate or in combination with a co-surfactant, preferably a C<sub>2</sub>-C<sub>10</sub> alkyl sulphate and/or a C<sub>8</sub>-C<sub>22</sub> alkyl or aryl sulphonate like C<sub>8</sub>-C<sub>22</sub> benzene sulphonate and/or another alkyl alkoxy  
35 sulphate apart the 3 ethoxylated one. Indeed the presence of said alkyl (ethoxy) 3 sulphate alone or in combination with a cosurfactant provides the desired




viscosity to the present compositions and delivers excellent stain removal properties to said compositions especially when used in any laundry application.

Typically such compositions according to the present invention have a viscosity  
5 between 25 cps and 1500 cps, preferably between 50 cps and 1100 cps, depending from the ratio between AE3S and the co-surfactant, when measured with a rheometer like carri-med CSL2-100® at the following viscosity parameters : angle : 1°58, gap : 60, diameter : 4.0 cm, iner : 63.60 at a temperature of 25°C and a shear rate of 30 1/sec.

10 Other anionic surfactants useful for deterative purposes can also be used herein. These can include salts (including, for example, sodium, potassium, ammonium, and substituted ammonium salts such as mono-, di- and triethanolamine salts) of soap, C<sub>9</sub>-C<sub>20</sub> linear alkylbenzenesulfonates, C<sub>8</sub>-C<sub>22</sub> primary or secondary  
15 alkanesulfonates, C<sub>8</sub>-C<sub>24</sub> olefinsulfonates, sulfonated polycarboxylic acids prepared by sulfonation of the pyrolyzed product of alkaline earth metal citrates, e.g., as described in British patent specification No. 1,082,179, C<sub>8</sub>-C<sub>24</sub> alkylpolyglycoethersulfates (containing up to 10 moles of ethylene oxide); alkyl ester sulfonates such as C<sub>14</sub>-C<sub>16</sub> methyl ester sulfonates; acyl glycerol  
20 sulfonates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, paraffin sulfonates, alkyl phosphates, isethionates such as the acyl isethionates, N-acyl taurates, alkyl succinamates and sulfosuccinates, monoesters of sulfosuccinate (especially saturated and unsaturated C<sub>12</sub>-C<sub>18</sub> monoesters) diesters of sulfosuccinate (especially saturated and unsaturated  
25 C<sub>6</sub>-C<sub>14</sub> diesters), sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic, nonsulfated compounds being described below), branched primary alkyl sulfates, alkyl polyethoxy carboxylates such as those of the formula  $RO(CH_2CH_2O)_kCH_2COO-M^+$  wherein R is a C<sub>8</sub>-C<sub>22</sub> alkyl, k is an integer from 0 to 10, and M is a soluble salt-forming cation. Resin acids  
30 and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tall oil. Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in U.S. Patent 3,929,678, issued December 30, 1975,  
35 to Laughlin, et al. at Column 23, line 58 through Column 29, line 23 (herein incorporated by reference).

Other suitable anionic surfactants for use herein also include acyl sarcosinate or mixtures thereof, in its acid and/or salt form, preferably long chain acyl sarcosinates having the following formula:

5



wherein M is hydrogen or a cationic moiety and wherein R is an alkyl group of from 11 to 15 carbon atoms, preferably of from 11 to 13 carbon atoms. Preferred  
10 M are hydrogen and alkali metal salts, especially sodium and potassium. Said acyl sarcosinate surfactants are derived from natural fatty acids and the amino-acid sarcosine (N-methyl glycine). They are suitable to be used as aqueous solution of their salt or in their acidic form as powder. Being derivatives of natural fatty acids, said acyl sarcosinates are rapidly and completely  
15 biodegradable and have good skin compatibility.

Accordingly, suitable long chain acyl sarcosinates to be used herein include C<sub>12</sub> acyl sarcosinate (i.e., an acyl sarcosinate according to the above formula wherein M is hydrogen and R is an alkyl group of 11 carbon atoms) and C<sub>14</sub> acyl  
20 sarcosinate (i.e., an acyl sarcosinate according to the above formula wherein M is hydrogen and R is an alkyl group of 13 carbon atoms). C<sub>12</sub> acyl sarcosinate is commercially available, for example, as Hamposyl L-30<sup>®</sup> supplied by Hampshire. C<sub>14</sub> acyl sarcosinate is commercially available, for example, as Hamposyl M-30<sup>®</sup> supplied by Hampshire.


25

Suitable nonionic surfactants for use herein include a class of compounds which may be broadly defined as compounds produced by the condensation of alkylene oxide groups (hydrophilic in nature) with an organic hydrophobic compound, which may be branched or linear aliphatic (e.g. Guerbet or  
30 secondary alcohol) or alkyl aromatic in nature. The length of the hydrophilic or polyoxyalkylene radical which is condensed with any particular hydrophobic group can be readily adjusted to yield a water-soluble compound having the desired degree of balance between hydrophilic and hydrophobic elements. Accordingly suitable nonionic synthetic surfactants include :

- (i) The polyethylene oxide condensates of alkyl phenols, e.g., the condensation products of alkyl phenols having an alkyl group containing from 6 to 20 carbon atoms in either a straight chain or branched chain configuration, preferably from 8 to 12 carbon atoms, with ethylene oxide, the said ethylene oxide being present in amounts equal to 10 to 25 moles of ethylene oxide per mole of alkyl phenol. The alkyl substituent in such compounds may be derived from polymerized propylene, diisobutylene, octane, and nonane;
- (ii) Those derived from the condensation of ethylene oxide with the product resulting from the reaction of propylene oxide and ethylene diamine products which may be varied in composition depending upon the balance between the hydrophobic and hydrophilic elements which is desired. Examples are compounds containing from about 40% to about 80% polyoxyethylene by weight and having a molecular weight of from about 5000 to about 11000 resulting from the reaction of ethylene oxide groups with a hydrophobic base constituted of the reaction product of ethylene diamine and excess propylene oxide, said base having a molecular weight of the order of 2500 to 3000;
- (iii) The condensation product of aliphatic alcohols having from 6 to 22 carbon atoms, in either straight chain or branched chain configuration, preferably from 8 to 18 carbon atoms, with from 2 to 35 moles of ethylene oxide, preferably from 4 to 25 and more preferably from 5 to 18. Example of this type of material are a coconut alcohol ethylene oxide condensate having from 5 to 18 moles of ethylene oxide per mole of coconut alcohol, the coconut alcohol fraction having from 9 to 14 carbon atoms;
- (iv) Trialkyl amine oxides and trialkyl phosphine oxides wherein one alkyl group ranges from 10 to 18 carbon atoms and two alkyl groups range from 1 to 3 carbon atoms; specific example is tetradecyl dimethyl phosphine oxide;
- (v) The condensation products of ethylene oxide with a hydrophobic base formed by the condensation of propylene oxide with propylene glycol; The hydrophobic portion of these compounds will preferably have a molecular

weight of from about 1500 to about 1800 and will exhibit water insolubility. The addition of polyoxyethylene moieties to this hydrophobic portion tends to increase the water solubility of the molecule as a whole, and the liquid character of the product is retained up to the point where the polyoxyethylene content is about 50% of the total weight of the condensation product, which corresponds to condensation with up to about 40 moles of ethylene oxide.

Particularly suitable nonionic surfactants for use herein are capped nonionic ethoxylated surfactants according to the formula:



wherein  $R_1$  is a  $C_8$ - $C_{18}$  linear or branched alkyl or alkenyl group, aryl group, alkaryl group, preferably,  $R_1$  is a  $C_{10}$ - $C_{15}$  alkyl or alkenyl group, more preferably a  $C_{10}$ - $C_{15}$  alkyl group;


$R_2$  is a  $C_2$ - $C_{10}$  linear or branched alkyl group, preferably a  $C_4$  group;

$R_3$  is a  $C_1$ - $C_{10}$  alkyl or alkenyl group, preferably a  $C_1$ - $C_5$  alkyl group, and

$n$  is an integer ranging in the range of from 1 to 20, preferably from 1 to 10, more preferably from 1 to 5; or mixtures thereof.

These surfactants are commercially available from BASF under the trade name Plurafac®, from HOECHST under the trade name Genapol® or from ICI under the trade name Symperonic®. Preferred capped nonionic ethoxylated surfactant of the above formula are those commercially available under the tradename Genapol® L 2.5 NR from Hoechst, and Symperonic® LF/CS 1100 from ICI.


Other suitable nonionic surfactants for use herein include the amine oxides corresponding to the formula:



wherein  $R$  is a primary alkyl group containing 6-24 carbons, preferably 10-18 carbons, and wherein  $R'$  and  $R''$  are, each, independently, an alkyl group containing 1 to 6 carbon atoms. The arrow in the formula is a conventional representation of a semi-polar bond. The preferred amine oxides are those in


which the primary alkyl group has a straight chain in at least most of the molecules, generally at least 70%, preferably at least 90% of the molecules, and the amine oxides which are especially preferred are those in which R contains 10-18 carbons and R' and R" are both methyl. Exemplary of the preferred amine oxides are the N-hexyldimethylamine oxide, N-octyldimethylamine oxide, N-decyldimethylamine oxide, N-dodecyl dimethylamine oxide, N-tetradecyldimethylamine oxide, N-hexadecyl dimethylamine oxide, N-octadecyldimethylamine oxide, N-eicosyldimethylamine oxide, N-docosyldimethylamine oxide, N-tetracosyl dimethylamine oxide, the corresponding amine oxides in which one or both of the methyl groups are replaced with ethyl or 2-hydroxyethyl groups and mixtures thereof. A most preferred amine oxide for use herein is N-decyldimethylamine oxide.

Other suitable nonionic surfactants for the purpose of the invention are the phosphine or sulfoxide surfactants of formula :



wherein A is phosphorus or sulfur atom, R is a primary alkyl group containing 6-24 carbons, preferably 10-18 carbons, and wherein R' and R" are, each, independently selected from methyl, ethyl and 2-hydroxyethyl. The arrow in the formula is a conventional representation of a semi-polar bond.

Suitable zwitterionic detergents for use herein comprise the betaine and betaine-like detergents wherein the molecule contains both basic and acidic groups which form an inner salt giving the molecule both cationic and anionic hydrophilic groups over a broad range of pH values. Some common examples of these detergents are described in U.S. Pat. Nos. 2,082,275, 2,702,279 and 2,255,082, incorporated herein by reference. Preferred zwitterionic detergent compounds have the formula :  $R^2$



wherein R1 is an alkyl radical containing from 8 to 22 carbon atoms, R2 and R3 contain from 1 to 3 carbon atoms, R4 is an alkylene chain containing from 1 to

3 carbon atoms, X is selected from the group consisting of hydrogen and a hydroxyl radical, Y is selected from the group consisting of carboxyl and sulfonyl radicals and wherein the sum of R1, R2 and R3 radicals is from 14 to 24 carbon atoms.


5

Amphoteric and ampholytic detergents which can be either cationic or anionic depending upon the pH of the system are represented by detergents such as dodecylbeta-alanine, N-alkyltaurines such as the one prepared by reacting dodecylamine with sodium isethionate according to the teaching of U.S. Pat. No. 2,658,072, N-higher alkylaspartic acids such as those produced according to the teaching of U.S. Pat. No. 2,438,091, and the products sold under the trade name "Miranol", and described in U.S. Pat. No. 2,528,378, said patents being incorporated herein by reference. Additional synthetic detergents and listings of their commercial sources can be found in McCutcheon's Detergents and Emulsifiers, North American Ed. 1980, incorporated herein by reference.

10

Cationic surfactants suitable for use in compositions of the present invention are those having a long-chain hydrocarbonyl group. Examples of such cationic surfactants include the ammonium surfactants such as alkyltrimethylammonium halogenides, and those surfactants having the formula:

15



25


wherein R<sup>2</sup> is an alkyl or alkyl benzyl group having from 8 to 18 carbon atoms in the alkyl chain, each R<sup>3</sup> is selected from the group consisting of -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH(CH<sub>3</sub>)-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, and mixtures thereof; each R<sup>4</sup> is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub> alkyl, benzyl ring structures formed by joining the two R<sup>4</sup> groups, and hydrogen when y is not 0; R<sup>5</sup> is the same as R<sup>4</sup> or is an alkyl chain wherein the total number of carbon atoms of R<sup>2</sup> plus R<sup>5</sup> is not more than about 18; each y is from 0 to about 10 and the sum of the y values is from 0 to about 15; and X is any compatible anion.

30

Quaternary ammonium surfactant suitable for the present invention has the formula (I):

35

27




Formula I

whereby R1 is a short chainlength alkyl (C6-C10)  
y is 2-4, preferably 3.

- 5 whereby R2 is H or a C1-C3 alkyl,  
whereby x is 0-4, preferably 0-2, most preferably 0,  
whereby R3, R4 and R5 are either the same or different and can be either a  
short chain alkyl (C1-C3) or alkoxyated alkyl of the formula II,  
10 whereby X<sup>-</sup> is a counterion, preferably a halide, e.g. chloride or methylsulfate.

78



Formula II

- 15 R6 is C1-C4 and z is 1 or 2.

Preferred quat ammonium surfactants are those as defined in formula I  
whereby

- R1 is C8, C10 or mixtures thereof, x=0,  
20 R3, R4, R5 = CH3.

Highly preferred cationic surfactants are the water-soluble quaternary  
ammonium compounds useful in the present composition having the formula :

- 25  $R_1R_2R_3R_4N^+X^-$  (i)

wherein R1 is C8-C16 alkyl, each of R2, R3 and R4 is independently C1-C4  
alkyl, benzyl, and  $-(C_2H_4O)_xH$  where x has a value from 2 to 5, and X is an  
anion. Not more than one of R2, R3 or R4 should be benzyl.

- 30 The preferred alkyl chain length for R1 is C12-C15 particularly where the alkyl  
group is a mixture of chain lengths derived from coconut or palm kernel fat or is  
derived synthetically by olefin build up or OXO alcohols synthesis. Preferred

groups for  $R_2R_3$  and  $R_4$  are methyl and the anion X may be selected from halide, methosulphate, acetate and phosphate ions.

Examples of suitable quaternary ammonium compounds of formulae (i) for use herein are :

- 5 coconut trimethyl ammonium chloride or bromide;  
decyl triethyl ammonium chloride;  
C<sub>12-15</sub> trimethyl ammonium chloride or bromide;  
myristyl trimethyl ammonium methyl sulphate;  
lauryl dimethyl benzyl ammonium chloride or bromide;  
10 lauryl dimethyl (ethenoxy)<sub>4</sub> ammonium chloride or bromide;  
choline esters (compounds of formula (i) wherein  $R_1$  is  
CH<sub>2</sub>-CH<sub>2</sub>-O-C-C<sub>12-14</sub> alkyl and  $R_2R_3R_4$  are methyl).

||

O

- 15 di-alkyl imidazolines

Other cationic surfactants useful herein are also described in U.S. Patent 4,228,044, Cambre, issued October 14, 1980 and in European Patent Application EP 000,224.

20

#### Optional polymers

- An optional component of the compositions herein is a polymer. Suitable polymers for use herein comprise monomeric units selected from the group consisting of unsaturated carboxylic acids, polycarboxylic acids, sulphonic acids, phosphonic acids and mixtures thereof. Co-polymerisation of the above monomeric units among them or with other co-monomers such as styrenesulfonic acid is also suitable.

- 30 Preferred examples of such polymers are the polymers and co-polymers of monomeric units selected from the group consisting of acrylic acid, maleic acid, vinylsulphonic acid and mixtures thereof. Also suitable for use herein are the above mentioned polymers and co-polymers which are modified in order to contain other functional groups such as aminophosphonic and/or phosphonic units. More preferred polymers are selected from the group consisting of  
35 polyacrylate polymers, co-polymers of acrylic and maleic acid, co-polymers of



styrene sulphonic acid and maleic acid, and mixtures thereof, preferably modified with aminophosphonic and/or phosphonic groups.

5 The molecular weight for these polymers and co-polymers is preferably below 100,000, most preferably between 500 and 50,000. Most suitable polymers and co-polymers for use herein will be soluble in an amount up to 0.1% by weight, in an aqueous composition comprising 5% by weight of sodium hypochlorite with its pH adjusted to 13 with sodium hydroxide.

10 Commercially available such polymers, suitable for use herein, are the polyacrylate polymers sold under the tradename Good-Rite® from BF Goodrich, Acrysol® from Rohm & Haas, Sokalan® from BASF, Norasol® from Norso Haas. Also suitable for use herein are the co-polymers of styrene sulphonic acid and maleic acid, commercially available under the tradename  
15 Versaflex® from National Starch such as Versaflex 157, as well as Acumer® terpolymers from Rohm and Haas, in particular Acumer® 3100. Preferred commercially available polymers are the polyacrylate polymers, especially the Norasol® polyacrylate polymers and more preferred are the polyacrylate polymer Norasol® 410N (MW 10,000) and the polyacrylate polymer modified  
20 with phosphonic groups Norasol® 440N (MW 4000) and its corresponding acid form Norasol® QR 784 (MW 4000).

A preferred polymer for use herein is a polyacrylate polymer modified with phosphonic groups commercially available under the tradename Norasol®  
25 440N (MW 4000) and its corresponding acid form Norasol® QR 784 (MW 4000) from Norso-Haas.

Mixtures of polymers as herein described may also be used in the present invention.

30

Polymers herein are preferably present in low amounts, i.e. in amounts of up to 10%, preferably up to 1%, more preferably up to 0.5% by weight, even more preferably from 0.001% to 0.3% by weight, and most preferably from 0.005% to 0.2% by weight of the composition.

Method of bleaching fabrics:

Particularly preferred surfaces to be bleached with the compositions herein are fabrics including for example clothes, curtains, drapes, bed linens, bath linens, table cloths, sleeping bags, tents, upholstered furniture and the like, and carpets.

Thus, the present invention further encompasses a method of bleaching fabrics which comprises the step of contacting said fabrics with a bleaching composition according to the present invention. In a preferred embodiment, the compositions used in said method of bleaching fabrics are liquid hypochlorite-containing compositions that may further comprise a chelating agent and/or a pH buffering component as defined hereinbefore. Said method according to the present invention delivers effective whiteness performance upon ageing of the compositions.

The compositions according to the present invention are applied to fabrics in a liquid form. Indeed, by "in a liquid form", it is meant herein the liquid compositions according to the present invention per se in neat or diluted form.

The compositions according to the present invention are typically used in diluted form in a laundry operation. By "in diluted form", it is meant herein that the compositions for the bleaching of fabrics according to the present invention may be diluted by the user, preferably with water. Such dilution may occur for instance in hand laundry applications as well as by other means such as in a washing machine. Said compositions can be diluted up to 500 times, preferably from 5 to 200 times and more preferably from 10 to 80 times.


More specifically, the method of bleaching fabrics according to the present invention comprises the steps of first contacting said fabrics with a bleaching composition according to the present invention, in its diluted form, then allowing said fabrics to remain in contact with said composition, for a period of time sufficient to bleach said fabrics, typically 1 to 60 minutes, preferably 5 to 30 minutes, then rinsing said fabrics with water. If said fabrics are to be washed, i.e., with a conventional composition comprising at least one surface active agent, said washing may be conducted together with the bleaching of said fabrics by contacting said fabrics at the same time with a bleaching composition

according to the present invention and said detergent composition, or said washing may be conducted before or after that said fabrics have been bleached.

- 5 In another embodiment of the present invention the method of bleaching fabrics comprises the step of contacting fabrics with a liquid bleaching composition according to the present invention, in its neat form, and allowing said fabrics to remain in contact with said bleaching composition for a period of time sufficient to bleach said fabrics, typically 5 seconds to 30 minutes, preferably 1 minute to  
10 10 minutes before rinsing said fabrics with water. If said fabrics are to be washed, i.e., with a conventional composition comprising at least one surface active agent, said washing may be conducted before or after that the fabrics have been bleached with the composition herein. In the embodiment of the present invention wherein the liquid bleaching composition of the present  
15 invention, is applied to the fabrics in its neat form, it is preferred that the level of hypohalite bleach, is from 0.01% to 5%, preferably from 0.1% to 3.5%, more preferably from 0.2% to 2% and most preferably from 0.2% to 1%.

- It is preferred to perform the bleaching methods herein before said fabrics are washed. Indeed, it has been observed that bleaching said fabrics with the  
20 compositions according to the present invention (diluted and/or neat bleaching methods) prior to washing them with a detergent composition provides superior whiteness and stain removal with less energy and detergent than if said fabrics are washed first, then bleached.


- 25 Alternatively instead of following the neat bleaching method as described herein above (pretreater application) by a rinsing step with water and/or a conventional washing step with a liquid or powder conventional detergent, the bleaching pre-treatment operation may also be followed by the diluted  
30 bleaching method as described herein before either in bucket (hand operation) or in a washing machine.



Claims


1. A liquid cleaning composition comprising an oxidising agent and a radical scavenger which is selected from the group consisting of:

(i)




wherein S is a solubilising group ( $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ), each X, Y, Z, W, are  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl groups, J is  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl group and M is H or an alkali metal, or mixtures thereof.

(ii)



naphtalene derivatives where in the position indicated with the numbers from 1 to 8 substituents are present (S, A, B, C, D, E, F, G, any substituents can occupy any position) wherein S is a solubilising group ( $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ) A, B, C, D, are  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl groups, E, F and G can be  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl group and M is H or an alkali metal, or mixtures thereof.


(iii)



Where wherein S is a solubilising group ( $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ), R' is C2-20 linear or branched alkyl chain; M is as above, P and N can be  $-\text{OR}'$  or  $-\text{H}$ ,  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OCH}_3$ , or a  $\text{C}_1$  to  $\text{C}_{10}$  primary and secondary alkyl groups.

(iv) and mixtures thereof.


- 2- A composition according to any preceding claim wherein the oxidising agent is a hypohalite, preferably hypochlorite bleach.
- 3- A composition according to any preceding claim additionally comprising a surfactant selected from the group consisting of anionic, nonionic, cationic, amphoteric, zwitterionic surfactants and mixtures thereof.
- 4- A composition according to claim 8 comprising an anionic surfactant selected from alkyl sulphate, alkyl ether sulphate and mixtures thereof.
- 5- A composition according to any preceding claim in thickened liquid, preferably aqueous form.
- 6- The use of a compound having the formula



wherein S is a solubilising group ( $-\text{COO}-\text{M}^+$  or  $-\text{SO}_3-\text{M}^+$ ), each X, Y, Z, W, are  $-\text{COO}-\text{M}^+$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{SO}_3-\text{M}^+$ ,  $-\text{NO}_2$ ,  $-\text{OR}$  (with R=linear or branched alkyl chain


C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl groups, J is -H, -COO-M<sup>+</sup>, -Cl, -Br, -SO<sub>3</sub>-M<sup>+</sup>, -NO<sub>2</sub>, -OR (with R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl group and M is H or an alkali metal, or mixtures thereof.

7- The use of a compound according to the formula



naphtalene derivatives where in the position indicated with the numbers from 1 to 8 substituents are present (S, A, B, C, D, E, F, G, any substituents can occupy any position) wherein S is a solubilising group (-COO-M<sup>+</sup> or -SO<sub>3</sub>-M<sup>+</sup>) A, B, C, D, are -COO-M<sup>+</sup>, -Cl, -Br, -SO<sub>3</sub>-M<sup>+</sup>, -NO<sub>2</sub>, -OR (with R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl groups, E, F and G can be -H, -COO-M<sup>+</sup>, -Cl, -Br, -SO<sub>3</sub>-M<sup>+</sup>, -NO<sub>2</sub>, -OR (with R=linear or branched alkyl chain C1-C20), or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl group and M is H or an alkali metal, or mixtures thereof as a radical scavenger.

8- The use of a compound having formula



Where wherein S is a solubilising group (-COO-M<sup>+</sup> or -SO<sub>3</sub>-M<sup>+</sup>), R' is C2-20 linear or branched alkyl chain; M is as above, P and N can be -OR' or -H, -COO-M<sup>+</sup>, -Cl, -Br, -SO<sub>3</sub>-M<sup>+</sup>, -NO<sub>2</sub>, -OCH<sub>3</sub>, or a C<sub>1</sub> to C<sub>10</sub> primary and secondary alkyl groups as a radical scavenger.





Abstract

The present invention relates to cleaning composition which comprise radical scavenger, for effective cleaning and improved storage stability.

